

ON STRICT SUNS IN $\ell^\infty(3)$

A. R. ALIMOV

A subset M of a normed linear space X is said to be a *strict sun* if, for every point $x \in X \setminus M$, the set of its nearest points from M is non-empty and if $y \in M$ is a nearest point from M to x , then y is a nearest point from M to all points from the ray $\{\lambda x + (1 - \lambda)y \mid \lambda > 0\}$. In the paper there obtained a geometrical characterisation of strict suns in $\ell^\infty(3)$. In comparison with [1] we establish a more precise property of strict suns.

A subset M of a normed linear space X is said to be a *strict sun* if, for every $x \in X \setminus M$, the set of its nearest points from M is non-empty and if $y \in M$ is the nearest point to x , then y is the nearest point from M to every point from the ray starting at y and passing through x . In this paper we obtain a geometric characterisation of strict suns in the space $\ell^\infty(3)$ (Theorem 1). This characterisation provides a more precise property of strict suns in $\ell^\infty(3)$ than Theorem A (see below) from [1] does.

1. Definitions and notation.

We will consider only real spaces.

Following K. Menger [2] and H. Berens and L. Hetzelt [3], a set $M \subset \mathbb{R}^n$ is called ℓ^1 -convex if, for all $x, y \in M$, $x \neq y$, there is a point $z \in M$, $z \neq x$, $z \neq y$, such that $\|x - y\|_{\ell^1} = \|x - z\|_{\ell^1} + \|z - y\|_{\ell^1}$ (here $\|\cdot\|_{\ell^1}$ is the standard ℓ^1 -norm on $\mathbb{R}^n$).

As it is shown in [3], ℓ^1 -convexity (together with closedness) proved to be a characteristic property of suns $\ell^\infty(n)$. (Here we recall that a set $M \subset X$ is a *sun* if, for every point $x \in X \setminus M$ there is a point y , that is nearest from M to x and such that y is a nearest point to all points from the ray starting at y and passing through x ; such a point y is called a *solar* point for x). It is clear that every strict sun is a sun. The inverse statement is not true in general. On suns, strict suns and other approximative sets see [4], [5].

An elegant characterisation of suns in $\ell^\infty(n)$ was obtained by H. Berens and L. Hetzelt [3] (see also [6]).

Theorem (H. BERENS, L. HETZELT). *A closed nonvoid set $M \subset \mathbb{R}^n$ is a sun in $\ell^\infty(n)$ if and only if it is ℓ^1 -convex.*

The first characterisation of strict suns in $\ell^\infty(n)$ in geometrical terms was independently obtained by B. Brosowski [7] and Ch. Dunham [8]. In [1] (Theorem A) there obtained a characterisation of strict suns in $\ell^\infty(n)$, which is similar to the metric characterisation due to Berens and Hetzelt.

1991 *Mathematics Subject Classification.* 41A65.

Key words and phrases. Strict sun; sun; $\ell^\infty(n)$; characterisation; best approximation; metrical convexity; ℓ^1 -convexity.

Supported by the Russian Fund for Basic Researches, Project No. 02-01-00248.

Here some necessary definitions are required.

Given a set $M \subset \mathbb{R}^n$, we define

$$\text{eqc}(M) = \text{eqc}_{\mathbb{R}^n}(M) := \{j \in \{1, \dots, n\} \mid x_j = y_j \text{ for all } x, y \in M\},$$

$$x = (x_1, \dots, x_n), \quad y = (y_1, \dots, y_n);$$

the abbreviation “eqc” stands for ‘equal coordinates’.

For $x, y \in \mathbb{R}^n$ and $M \subset \mathbb{R}^n$, we define

$$x \not\vdash_M y, \text{ if } x_j \neq y_j \text{ for all } j \notin \text{eqc}(M);$$

$$x \not\vdash y, \text{ if } x_i \neq y_i \text{ for all } i = 1, \dots, n.$$

A set M is said to be *strictly ℓ^1 -convex* if

- 1) it is ℓ^1 -convex, and
- 2) for every $x, y \in M$, $x \not\vdash_M y$, there is an ℓ^1 -geodesic segment $k(t) \subset M$, $0 \leq t \leq 1$, connecting x and y for which the functions $[0, 1] \ni t \mapsto k_j(t)$ are monotonic on t for $j = 1, 2, \dots, n$ and strictly monotonic for $j \notin \text{eqc}(M)$.

In our terminology, the ℓ^1 -convexity of M means [3] that any two points $x, y \in M$ can be connected by an ℓ^1 -geodesic segment $k(\cdot) \subset M$, with monotonic in t coordinate functions $k_i(t)$ (which are not necessarily strictly monotonic).

Following [1], by a *main cocross* in $\mathbb{R}^n$ we will understand the set

$$c^n(x) = \{y \in \mathbb{R}^n \mid \text{card eqc}(y, x) \geq 1\}, \quad x \in \mathbb{R}^n \text{ is fixed,}$$

and by a *cocross* $c_J(x)$ with respect to the assembly of coordinates $J = \{j_1, \dots, j_k\} \subset \{1, \dots, n\}$ we will understand the set

$$c_J(x) = c_J^n(x) = \left\{ y \in \mathbb{R}^n \mid \begin{array}{ll} y_j = \text{const}(j) & \text{for } j \in J, \\ \text{card}(\text{eqc}(x, y) \cap J) \geq 1 & \text{for } i \notin J. \end{array} \right\}$$

In addition, to exclude degenerate cases, we will *always* assume that $k \leq n - 2$; i.e., $\text{card } J \leq n - 2$ in the definition of $c_J(x)$. Finally, a *cocross* is any set $C \subset \mathbb{R}^n$ such that $C \subset c_{\text{eqc}(C)}(x)$ for some $x \in \mathbb{R}^n$. (Note that always $\text{card eqc}(C) < n - 1$.)

For $x \in \mathbb{R}^n$ we will define

$$\text{cr}(x) = \text{cr}^n(x) = \{y \in \mathbb{R}^n \mid \text{card eqc}(x, y) \geq n - 1\} \text{ a } \textit{main cross} \text{ in } \mathbb{R}^n.$$

A *cross* C is a subset of some main cross in $\mathbb{R}^n$ such that $\text{eqc}(C) = \emptyset$; we will call it $\mathbb{R}^n$ -*cross*. It is clear that any $\mathbb{R}^n$ -cross is a cocross.

The next theorem [1] characterises strict suns in $\ell^\infty(n)$.

Theorem A. *Let $\emptyset \neq M \subset \mathbb{R}^n$ be closed. Then M is a strict sun in $\ell^\infty(n)$ if and only if it is 1) strictly ℓ^1 -convex and 2) not a cocross.*

Conditions 1) and 2) are not independent ones. In [1] it was mentioned that $\mathbb{R}^3$ -cross $\text{cr}^3(x)$ is an ℓ^1 -convex set. However, the main cocross $c^n(x)$ is, clearly, not ℓ^1 -convex.

2. The main theorem.

Here we prove a result in which the characteristic conditions in Theorem A are rectified for $n \leq 3$. Namely, condition 2) is replaced with the more strong one: *a set is not an $\mathbb{R}^3$ -cross*. We also prove Proposition 1, which characterises strict ℓ^1 -convex cocrosses in $\mathbb{R}^3$ among all cocrosses in $\mathbb{R}^3$.

Theorem 1. *Let $\emptyset \neq M \subset \mathbb{R}^3$ be closed. Then M is a strict sun in $\ell^\infty(3)$ if and only if it is 1) strictly ℓ^1 -convex and 2) not an $\mathbb{R}^3$ -cross.*

In other words, for a closed set M its strict ℓ^1 -convexity is necessary for M to be a sun in $\ell^\infty(n)$, and is sufficient provided that M is not a cross (for $\ell^\infty(3)$) or not a cocross (for $\ell^\infty(n)$).

In the plane $\mathbb{R}^2$ the situation is trivial: any cocross in $\mathbb{R}^2$ is an $\mathbb{R}^2$ -cross, which evidently is not a strict ℓ^1 -convex set. It follows that in $\mathbb{R}^2$ Theorem A coincides with Theorem 1.

The next proposition characterises strict ℓ^1 -convex cocrosses among all cocrosses in $\mathbb{R}^3$. We will prove it later.

Proposition 1. *A cocross in $\mathbb{R}^3$ is strictly ℓ^1 -convex if and only if it is a connected $\mathbb{R}^3$ -cross.*

(Topological dimension of an $\mathbb{R}^n$ -cross K is always 1, therefore its connectedness is equivalent to arcwise connectedness and, because of the inclusion $K \subset \text{cr}(x)$, is equivalent to ℓ^1 -convexity.)

3. Preliminary results.

As usual, by e_1, e_2, e_3 we denote the standard basis of unit vectors in $\mathbb{R}^3$.

Lemma 1. *Let $\emptyset \neq M \subset \mathbb{R}^3$ has the following properties:*

- 1) $0, e_1, e_2, e_3 \in M$,
- 2) M is not a cross;
- 3) *any two points $x, y \in M$ such that $x \neq y$ can be connected by a curve $k(t) \subset M$, $k(0) = x$, $k(1) = y$, having strictly monotonic coordinate functions.*

Then there is a point $w \in M$ such that $w \neq 0$.

Proof. Let $x \in M$ and let $x \notin \text{cr}(0)$. Such a point exists by the second assumption. If $x \neq 0$ then the desired statement is proved. Suppose that $x \in c^3(0)$. Without loss of generality assume that $x_1 = 0$. Since $M \not\subset \text{cr}(0)$ then $x \neq e_1$. From our assumptions it follows that x and e_1 are connected by the curve $k(t) \subset M$, $k(0) = x$, $k(1) = e_1$, with strictly monotonic in t coordinate functions. It is clear that $c^3(0) \subset c^3(e_1) \cup c^3(x)$. Now the strict monotonicity of functions $k_i(t)$, $i = 1, 2, 3$, implies that $k(t) \notin c^3(e_1) \cup c^3(x)$ for $0 < t < 1$, and therefore $k(t) \notin c^3(0)$; i.e., $k(t) \neq 0$ for every $t \in (0, 1)$. This proves the lemma, since $k(t) \subset M$. $\square$

For $x \in X$, $P_M x$ denotes the set of all nearest points from M to X ; i.e.,

$$P_M x = \{y \in M \mid \|x - y\| = \rho(x, M) := \inf_{z \in M} \|x - z\|\}.$$

The next evident lemma is well known (see e.g. [9]).

Lemma 2. *Let X be a normed linear space and let $M \subset X$, $x \in X$, $x \notin M$. Suppose that $\hat{y} \in M$ is a solar point from M to x and $y \in P_M x$. Then $[y, \hat{y}] \subset S(x, \|x - y\|)$.*

The next result will be used in proof of Proposition 1.

Lemma 3. *Suppose that a set $M \subset \mathbb{R}^3$ is ℓ^1 -convex and is not a cross. Then there exist points $x, y \in M$ such that $x \neq y$.*

Proof. Suppose that the statement is not true. Without loss of generality assume that $0 \in M$ and $\text{eqc}(M) = \emptyset$. Since $M \not\subset \text{cr}(0)$, there exists a point $x \in M$ such

that $x \in c^3(0)$, $x \notin \text{cr}(0)$. (Note that the case $M \not\subset c^3(0)$ immediately implies the desired statement.)

Without loss of generality we assume $x = (x_1, x_2, 0)$, $x_1, x_2 \neq 0$. Since $\text{eqc}(M) = \emptyset$, there exists a point $y \in M$ such that $y_3 \neq 0$. By the assumption, $\text{eqc}(x, y) \neq \emptyset$, therefore either $y_1 = x_1$, $y_2 = 0$, or $y_2 = x_2$, $y_1 = 0$. Without loss of generality we assume that the second case is fulfilled; i.e., $y = (0, x_2, y_3)$. Denote $\xi = (0, x_2, 0)$. From ℓ^1 -convexity of M it follows that $[y, \xi] \cup [\xi, x] \subset M$. Again applying ℓ^1 -convexity, we have $[0, \xi] \subset M$.

Now we have $\xi, 0, x, y \in M$. After the appropriate translation of the origin into the point ξ and the appropriate combination of orthogonal pivots and extensions, the points $x, y, 0$ will become e_1, e_2, e_3 in the new coordinate system, the point ξ will be its origin, and the set M will be represented as $\tilde{M}$, having the same properties as M had. By Lemma 1, there is a point $\tilde{w} \in \tilde{M}$, $\tilde{w} \neq 0$. The point $\tilde{w}$ in the original system will become w , for which will be $w \neq \xi$. Now the desired pair (w, ξ) , $w \neq \xi$ of points is found. Lemma 3 is proved. $\square$

4. Proof of Theorem 1.

In [1] it is proved that a strict sun in $\ell^\infty(n)$ is strictly ℓ^1 -convex and is not a cocross. Since a cross is certainly a cocross, we proved the “ONLY IF” part in the theorem.

“IF”. It is enough [1] to consider the following setting: $\text{eqc}(M) = \emptyset$.

By Berens–Hetzelt theorem the set M is a sun. We will prove that M is a strict sun by applying the well-known lemma of E. B. Oshman and A. Brøndsted (see e.g. [4, Chapter 3]); in accordance with this result it will be enough to prove that

$$\forall x \notin M \text{ and } \forall y \in P_M x \quad \mathring{K}(y, x) \cap M = \emptyset, \quad (1)$$

where

$$\mathring{K}(y, x) = \{z \mid [z, y] \cap \mathring{B}(x, \|x - y\|) \neq \emptyset\}$$

is the open supporting cone for the ball $B(x, \|x - y\|)$ at the point y (equivalent definitions of supporting cones one can find in [4, Chapter 3]).

Without loss of generality we assume $x = 0$, $\rho(0, M) = \|y\| = 1$, and $y_i \geq 0$, $i = 1, 2, 3$. Let us denote $J = \{j \mid y_j = 1\}$. Since $\|y\| = 1$ and $y_i \geq 0$, we have $J \neq \emptyset$.

Suppose that (1) is false; i.e., $\mathring{K}(y, 0) \cap M \neq \emptyset$.

Let $H = c^3(y) \cap \mathring{K}(y, 0)$.

1. Suppose at first that $w \in \mathring{K}(y, 0) \cap M$ and $w \notin H$. It follows that $w \notin c^3(y)$. Therefore $w \neq y$, and, by the strict ℓ^1 -convexity of M , it implies that w and y can be connected by a strictly monotonic ℓ^1 -geodesic segment $k(t) \subset M$, $k(0) = y$, $k(1) = w$.

Let us prove that $k(t) \subset \mathring{B}(0, 1)$ for all sufficiently small t . Under our assumptions we have $\mathring{K}(y, 0) = \{z \mid z_j < 1 \text{ for } j \in J\}$. Since $w \in \mathring{K}(y, 0)$ and since coordinate functions of the curve $k(t)$ are strictly monotonic, we have $w_j < k_j(t) < 1$ for $j \in J$ and $0 < t < 1$.

Since $k_j(t) \rightarrow y_j = 1$ for $t \rightarrow 0$ and $j \in J$, it follows that for some $\varepsilon > 0$

$$0 < k_j(t) < 1 \quad j \in J \quad t \in (0, \varepsilon). \quad (2)$$

Further, $k_m(t) \rightarrow y_m$ as $t \rightarrow 0$, $1 \leq m \leq 3$, and $0 \leq y_i < 1$ for $i \notin J$. Therefore, for some $\varepsilon_1 > 0$ we have

$$|k_i(t) - y_i| < 1 - y_i \quad i \notin J, \quad 0 < t < \varepsilon_1.$$

From this and from (2) it follows that

$$-1 < k_m(t) < 1 \quad t \in (0, \min\{\varepsilon, \varepsilon_1\}), \quad m = 1, 2, 3.$$

Therefore $k(t) \subset \mathring{B}(0, 1)$ for $t \in (0, \min\{\varepsilon, \varepsilon_1\})$. But this is impossible, since $y \in P_M 0$.

2. Suppose now that $M \cap \mathring{K}(y, 0) \subset H$. Fix $u \in H \cap M$. Since M is strictly ℓ^1 -convex, by Lemma 3 there are points $\xi, \eta \in M$, $\xi \neq \eta$ which are connected by a curve $\varkappa(t) \subset M$, $\varkappa(0) = \xi$, $\varkappa(1) = \eta$ having strictly monotonic in t coordinate functions. This implies that there is a point $v = \varkappa(t_0) \in M$, $0 < t_0 < 1$, such that $v \neq u$. From strict ℓ^1 -convexity of M it follows that u and v can be connected by a strictly monotonic ℓ^1 -geodesic segment $k(t) \subset M$, $0 \leq t \leq 1$, $k(0) = u$, $k(1) = v$. Since $u \in \mathring{K}(y, 0)$, we have that $u_j < 1$ for $j \in J$ and therefore $k_j(t) < 1$ for $j \in J$, $0 < t < \varepsilon_2$ for some ε_2 ; i.e., $k(t) \in \mathring{K}(y, 0)$ for $0 < t < \varepsilon_2$. From strict monotonicity of $k(t)$ it follows that there exists $\varepsilon_3 > 0$ such that $k(t) \neq y$ for all $t \in (0, \varepsilon_3)$. Now we have that for $0 < t < \min\{\varepsilon_2, \varepsilon_3\}$ the curve $k(t) \subset M$ is contained in $\mathring{K}(y, 0)$ and is not contained in H , which contradicts the assumption $M \cap \mathring{K}(y, 0) = H$. Thus the assumption $\mathring{K}(y, 0) \cap M \neq \emptyset$ is false and by Oshman–Brøndsted’s lemma M is a strict sun. The theorem is proved. $\square$

Proof of Proposition 1. The “ONLY IF” part. Let $K \subset \mathbb{R}^3$ be an ℓ^1 -convex cocross which is not an $\mathbb{R}^3$ -cross. Without loss of generality we assume that $K \subset c^3(0)$.

Let us at first consider the case when $\text{eqc}(K) \neq \emptyset$. We can assume that $\{3\} \in \text{eqc}(K)$ and that $K \subset c^2(0)$. By the definition of a cocross $\text{card eqc}(K) < 2$, whence $\text{eqc}(K) = \{3\}$; i.e., $\text{eqc}_{\mathbb{R}^2}(K) = \emptyset$. It means that K has points of the form $x = (\xi, 0)$ and $y = (0, \eta)$, $\xi, \eta \neq 0$. It is clear that $x \neq_K y$. However, since $K \subset c^2(0)$, it follows that x and y cannot be connected by a curve $k(t) \subset K$ with strictly monotonic in t coordinate functions $k_i(t)$, $i = 1, 2$, which contradicts the strict ℓ^1 -convexity of K .

Now let us consider the case when $\text{eqc}(K) = \emptyset$. Since K is not a cross, by Lemma 3 there are $x, y \in K$ such that $x \neq y$. By the strict ℓ^1 -convexity of K such x and y are connected by a curve $k(t) \subset K$ with strictly monotonic in t coordinate functions $k_i(t)$, $i = 1, 2, 3$. It is easy to see that this is impossible, since $K \subset c_{\mathbb{R}^3}(0)$.

This contradiction shows that a strictly ℓ^1 -convex cocross K has to be an $\mathbb{R}^3$ -cross, which is ℓ^1 -convex, and therefore connected.

The “If” part. We will prove that every connected $\mathbb{R}^n$ -cocross K for $n \geq 3$ is strictly ℓ^1 -convex. We may assume $K \subset c^n(0)$. By definition, $\text{eqc}(K) = \emptyset$. Every point from K is of the form: $(0, \dots, \xi, 0, \dots, 0)$. Since $n \geq 3$, we see that every two points from K have at least one common coordinate; i.e., $x \neq y$ is never fulfilled for $x, y \in K$. $\square$

Remark. In $\mathbb{R}^4$ there exists a strictly ℓ^1 -convex cocross which is not a cross. In fact, let us consider the set

$$K = \left\{ x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \left| \begin{array}{ll} x_1 = x_2 = x_3 = 0, & \text{if } x_4 \neq 0 \\ \text{card eqc}(x, 0) \geq 1, & \text{if } x_4 = 0 \end{array} \right. \right\}.$$

(The set K is union of the main cocross $c^3(0)$ in $\mathbb{R}^3$ and of the line $\{x \in \mathbb{R}^4 \mid x_1 = x_2 = x_3 = 0\}$ which is orthogonal to $c^3(0)$ in $\mathbb{R}^4$.) Now $K \subset c^4(0)$ and since $\text{eqc}(K) = \emptyset$, K is a cocross. However, all two points from K have at least one common coordinate. Now the strict ℓ^1 -convexity of K is evident.

REFERENCES

1. A. R. Alimov, *Geometrical characterization of strict suns in $\ell^\infty(n)$* , Mathematical Notes **70** (2001), no. 1, 3–10.
2. K. Menger, *Untersuchungen über allgemeine Metrik*, Math. Ann. **100** (1928), 75–163.
3. H. Berens, L. Hetzelt, *Die Metrische Struktur der Sonnen in $\ell^\infty(n)$* , Aequat. Math. **27** (1984), 274–287.
4. L. P. Vlasov, *Approximative properties of sets in normed linear spaces*, Russian Math. Surveys **28** (1973), no. 6, 1–66.
5. M. I. Karlov, I. G. Tsar'kov, *Convexity and connectedness of Chebyshev sets and suns*, Fundam. Prikl. Mat. **3** (1997), no. 4, 967–978.
6. A. L. Brown, *Suns in normed linear spaces which are finite-dimensional*, Math. Ann. **279** (1987), 87–101.
7. B. Brosowski, *Nich-Lineare Tschebyscheff-Approximation.*, B. I. Hochschulschriften., vol. 808/808a, ■
Hochschultaschenbücher Verlag, Bibliographisches Institut, Mannheim, 1968.
8. Ch. B. Dunham, *Characterizability and uniqueness in real Chebyshev approximation*, J. Appr. Th. **2** (1969), 374–383.
9. B. Brosowski, F. Deutsch, *Radial continuity of set-valued metric projections*, J. Approximation Theory **11** (1974), 236–253.